

Column sequences of cellular automata

Eric Rowland



Laboratoire de Combinatoire et d'Informatique Mathématique,
Université du Québec à Montréal

October 24, 2012


- 1 Cellular automata
- 2 Automatic sequences
- 3 Columns of linear cellular automata







One-dimensional cellular automata

- alphabet Σ of size k (for example $\{0, 1, \dots, k - 1\}$)
- function $i : \mathbb{Z} \rightarrow \Sigma$ (the initial condition)
- function $f : \Sigma^d \rightarrow \Sigma$ (the local update rule)




Naming scheme: $11111010_2 = 250$.






Binomial coefficients

Binomial coefficients modulo k are produced by cellular automata.



The local rule is $f(x, y, z) = x + z \text{ modulo } k$.


Boundary growth



Column sequences

characteristic sequence of 2^n

bits of π




characteristic sequence of n^2

statistically random sequences

Finiteness condition


The initial condition is eventually periodic in both directions.

Coarse-graining reduces to constant background.



Characteristic sequence of primes

A 16-color rule depending on 3 cells that computes the primes:



- 1 Cellular automata
- 2 Automatic sequences
- 3 Columns of linear cellular automata


The Thue–Morse sequence

$$T(n) = \begin{cases} 0 & \text{if the binary representation of } n \text{ has an even number of 1s} \\ 1 & \text{if the binary representation of } n \text{ has an odd number of 1s.} \end{cases}$$

The Thue–Morse sequence $T(n)_{n \geq 0}$ is

01101001100101101001011001101001

$T(n)$ occurs as a column of this $d = 5$ automaton:



Thue–Morse fun facts

- A **cube** is a word of the form www where w is a nonempty word. The infinite Thue–Morse word is cube-free:

01101001100101101001011001101001...

- Multigrades:

$$0^0 + 3^0 + 5^0 + 6^0 = 1^0 + 2^0 + 4^0 + 7^0 = 4$$

$$0^1 + 3^1 + 5^1 + 6^1 = 1^1 + 2^1 + 4^1 + 7^1 = 14$$

$$0^2 + 3^2 + 5^2 + 6^2 = 1^2 + 2^2 + 4^2 + 7^2 = 70$$


In general,
$$\sum_{n=0}^{2^\ell-1} (-1)^{T(n)} n^m = 0 \quad \text{for } 0 \leq m \leq \ell - 1.$$

- Interesting products:
$$\prod_{n \geq 0} \left(\frac{2n+2}{2n+1} \right)^{(-1)^{T(n)}} = \sqrt{2}$$

k -automatic sequences

The Thue–Morse sequence is 2-automatic.


It is computed by a **deterministic finite automaton with output** in base 2:



A sequence $s(n)_{n \geq 0}$ is **k -automatic** if there is k -DFAO whose output is $s(n)$ when fed the base- k digits of n .

Some 2-automatic sequences

- The characteristic sequence of powers of 2 is 2-automatic:




- The Rudin–Shapiro sequence

$$s(n) = \begin{cases} 0 & \text{if the binary representation of } n \text{ has an even number of 11s} \\ 1 & \text{if the binary representation of } n \text{ has an odd number of 11s} \end{cases}$$


is 2-automatic:

00010010000111010001001011100010 ...




Tower of Hanoi

- The minimal solution to the “infinite” tower of Hanoi puzzle is 2-automatic.



Dragon curve

- The sequence of folds in the paperfolding curve is 2-automatic.




Automatic sequences

Automatic sequences have been very well studied.

Büchi 1960:

If $s(n)$ is eventually periodic, then $s(n)$ is k -automatic for every $k \geq 2$.

Several natural generalizations of automatic sequences are known.



Christol's theorem

Let p be a prime.

Let \mathbb{F}_q be a finite field of characteristic p .

Theorem (Christol et al 1980)

A sequence $s(n)_{n \geq 0}$ of elements in \mathbb{F}_q is p -automatic if and only if the formal power series $\sum_{n \geq 0} s(n)t^n$ is algebraic over $\mathbb{F}_q(t)$.

The generating function $G(t) = \sum_{n \geq 0} T(n)t^n$ is algebraic over $\mathbb{F}_2(t)$:

$$tG(t) + (1 + t)G(t)^2 + (1 + t^4)G(t)^4 = 0.$$

The assumption $s(n) \in \mathbb{F}_q$ is not restrictive:

For a sequence on Σ , any injection $\Sigma \hookrightarrow \mathbb{F}_q$ gives an algebraic series.

- 1 Cellular automata
- 2 Automatic sequences
- 3 Columns of linear cellular automata**

Linear cellular automata

A cellular automaton is **linear** if the local rule $f : \mathbb{F}_q^d \rightarrow \mathbb{F}_q$ is \mathbb{F}_q -linear.

For example, the Pascal's triangle modulo p cellular automaton with $f(x, y, z) = x + z$ is linear.

Theorem (Litow–Dumas 1993)

Every column of a linear cellular automaton over \mathbb{F}_p is p -automatic.

The proof uses two theorems about formal power series — Christol's theorem and Furstenberg's theorem.

Furstenberg's theorem

The **diagonal** of a bivariate series $\sum_{n \geq 0} \sum_{m \geq 0} a(n, m)t^n x^m$ is

$$\sum_{n \geq 0} a(n, n)t^n.$$

Theorem (Furstenberg 1967)

A formal power series $G(t)$ is algebraic over $\mathbb{F}_q(t)$ if and only if $G(t)$ is the diagonal of a rational series $F(t, x)$.

Sketch of Litow–Dumas proof

Represent the n th row $\cdots a(n, -1) a(n, 0) a(n, 1) \cdots$ by

$$R_n(x) = \cdots + a(n, -1)x^{-1} + a(n, 0)x^0 + a(n, 1)x^1 + \cdots ,$$

which is rational since the initial condition is eventually periodic.

Linearity of the rule means $R_{n+1}(x) = C(x)R_n(x)$ for some $C(x)$.
For Pascal's triangle, $C(x) = x + \frac{1}{x}$.

Then the bivariate series $F(t, x) = \sum_{n \geq 0} \sum_{m \in \mathbb{Z}} a(n, m)t^n x^m = \sum_{n \geq 0} R_n(x)t^n = \sum_{n \geq 0} (C(x)t)^n R_0(x)$ is rational.

Column m of $F(t, x)$ is the diagonal of $x^{-m}F(tx, x)$, hence it is algebraic (Furstenberg) and hence p -automatic (Christol).

What about the converse?

We can reverse the proof, using the other directions of Christol's and Furstenberg's theorems.

Issue 1: We may not get a recurrence for $R_n(x)$ of order 1.

In general, $C_0(x)R_n(x) = \sum_{i=1}^d C_i(x)R_{n-i}(x)$.

To deal with this, we consider a cellular automaton with **memory**.

Issue 2: We need $C_0(x)$ to be a (nonzero) monomial so that each $\frac{C_i(x)}{C_0(x)}$ is a Laurent polynomial, so that the update rule is local.

Constructing a Thue–Morse cellular automaton

Christol's theorem gives that $x = \sum_{n \geq 0} T(n)t^n$ satisfies

$$tx + (1 + t)x^2 + (1 + t^4)x^4 = 0.$$

Replace $x \mapsto 0 + 1t + 1t^2 + t^2x$, and divide by t^3 .

Then $G(t) := \sum_{n \geq 0} T(n+3)t^n$ satisfies $P(t, G(t)) = 0$, where

$$P(t, x) = (t^2 + t^9) + x + (t + t^2)x^2 + (t^5 + t^9)x^4.$$

By Furstenberg's theorem, $T(n+2)$ is the coefficient of x^{-2} in $R_n(x)$:

$$\frac{P_x(t, x)}{P(t, x)} = \frac{1}{x} + t + \left(\frac{1}{x^2} + 1 + x \right) t^2 + \dots = \sum_{n \geq 0} R_n(x) t^n.$$

$R_n(x)$ satisfies the recurrence

$$R_n(x) = xR_{n-1}(x) + \left(\frac{1}{x} + x \right) R_{n-2}(x) + x^3 R_{n-5}(x) + \left(\frac{1}{x} + x^3 \right) R_{n-9}(x)$$

for all $n \geq 10$, which determines a linear cellular automaton rule with memory 9.

Restoring initial terms


Extend the memory to $9 + 3 = 12$ without introducing dependence on the earliest 3 rows.

Then $T(n)_{n \geq 0}$ occurs in Column -2 from initial conditions R_{-2}, \dots, R_9 .



0110100110010110 ...

Thue–Morse cellular automaton with memory 12




Theorem (Rowland–Yassawi 2012)

Every p -automatic sequence of elements in \mathbb{F}_q occurs as a column of a linear cellular automaton over \mathbb{F}_q with memory whose initial conditions are eventually periodic in both directions.

Combined with the Litow–Dumas result, we have a new characterization of p -automatic sequences (for prime p).


Rudin–Shapiro cellular automaton with memory 20



Baum–Sweet cellular automaton with memory 27

The Baum–Sweet sequence $1\ 1\ 0\ 1\ 1\ 0\ 0\ 1\ 0\ 1\ 0\ 0\ \dots$ is defined by

$$s(n) = \begin{cases} 0 & \text{if the binary representation of } n \\ & \text{contains a block of 0s of odd length} \\ 1 & \text{if not.} \end{cases}$$



If we give up linearity, we can get a cellular automaton without memory.

Corollary

Every p -automatic sequence occurs as a column of a cellular automaton whose initial condition is eventually periodic in both directions.


A cellular automaton rule is **invertible** if it has an inverse rule.

In other words, it can be evolved backward in time as well as forward.

Corollary

If $s(n)_{n \geq 0}$ is a p -automatic sequence, then for some $r \geq 0$ the sequence $s(n)_{n \geq r}$ occurs as a column of an invertible cellular automaton with memory.

Invertible Rudin–Shapiro cellular automaton



Open questions

- Given a p -automatic sequence on the alphabet $\Sigma \subset \mathbb{F}_q$, one can find a cellular automaton (without memory) with at most $q^{d+r+1} + |\Sigma|$ states containing the sequence. Can this bound be improved?
- Does there exist a 3-automatic sequence $s(n)_{n \geq 0}$ on a binary alphabet such that $s(n)$ is not eventually periodic and $s(n)$ occurs as a column of a (nonlinear) 2-state cellular automaton?
- Does every k -automatic sequence occur in a cellular automaton (if k is not prime)?
- Exhibit a sequence that does not occur as the column of a cellular automaton.